

Open Access Article

## CLINICAL, IMMUNOLOGICAL AND MEDICO-SOCIAL ASPECTS OF ALLERGIC DISEASES IN CHILDREN

**Ozoda Rajabturdievna Yomgurova**

Department of Pediatrics, Bukhara State Medical Institute, Bukhara, Uzbekistan

### Abstract

This article reveals the clinical and immunoregressive features of immunologic and allergic diseases in children, in addition to analyzing and analyzing the allergic characteristics of children in three developmental periods, the research demonstrates the scientific achievements of world scientists.

**Keywords:** Allergy, disimmunoglobulinemia, antigen, pregnant women, cytokine.

### 抽象的

本文揭示了儿童免疫学和过敏性疾病的临床和免疫退行特征，除了分析和分析了儿童三个发展时期的过敏特征外，该研究还展示了世界科学家的科学成就。

关键词：过敏，去免疫球蛋白血症，抗原，孕妇，细胞因子。

Today, the term "allergy" means an undesirable specific immune response, the implementation of various pathogenetic mechanisms, which can lead to allergic diseases [5, 9].

High antigenic loads significantly increase the risk of atopy formation: mother's smoking, toxicosis, irrational drug therapy of pregnant women, exposure to occupational allergens, unilateral carbohydrate nutrition, abuse of foods with obligate food allergens, etc. The exclusion of these moments is an important preventive factor [2, 12].

The links of the complex chain of an integrated approach to the treatment of allergic diseases, in addition to adequate basic therapy and the achievement of immunological tolerance, are preventive measures and the exclusion of the effect of triggers [5]. Immunoprophylaxis is undoubtedly one of the most important links in the complex therapy of patients with allergic diseases.

However, significant success can be achieved only by making up all the links of this important chain: to carry out fully and timely preventive measures and the elimination of trigger factors [6, 10].

Analyzing the literature data on the impact of allergens in rural children and the manifestation of allergic diseases, despite all the variety of treatment methods, the number of patients with allergic diseases is increasing every year all over the world. In addition, more and more children with immunological disorders are identified - dysimmunoglobulinemia, imbalance in the cytokine system, etc. Therefore, the correction of disorders in the immune system in such patients by selecting the most optimal methods of treatment is an urgent problem [1, 7].

Research in recent years has proven that allergy symptoms should not be considered the only marker of allergic disease [8]. Allergic

Received: April 12, 2021 / Revised: May 11, 2021 / Accepted: June 13, 2021 / Published: July 23, 2021

About the authors: Ozoda Rajabturdievna Yomgurova, Department of Pediatrics, Bukhara State Medical Institute, Bukhara, Uzbekistan

Corresponding author- Ozoda Rajabturdievna Yomgurova

inflammation and sensitization may not manifest clinically for a long time, but they are necessarily present, contributing to the progression of allergic diseases [9].

Of course, treatment tactics depend on the stage of the disease. If during the period of exacerbation the main therapeutic task is the elimination of acute allergic reactions, then during the period of remission it is to prevent progression and prevent relapse by changing the general reactivity and enhancing the ability to respond appropriately [10]. In addition, in patients with allergic diseases, the use of non-drug methods of therapy and rehabilitation is now becoming increasingly important, given their good combination with standard treatment regimens and the practical absence of side effects [12].

Dysfunction of the immune system is manifested by a decrease in the resistance of children to infectious agents, a decrease in the activity of the immune system. In the history of patients with a severe form of atopy, frequent exacerbations of the foci of the infectious process are observed. Immunodeficiencies in all parts of the immune system against the background of suppression in the system of T cell regulation in moderate to severe and severe course of the disease leads to the development of an autoimmune component [16].

In recent years, a significant role in the formation of the body's immune response has been assigned to the formation of the intestinal normal microflora. This process, which occurs in the first year of a child's life, has a long-term effect, laying down the characteristics of the body's immune response to external infectious and food antigens, predetermining the development of a particular pathology in the future. The process of recovery of the intestinal micro-biocenosis in a child is individual and depends on many factors: the state of health of the mother, the method of delivery,

the type of feeding, the use of antibiotics, etc. [14]. A decrease in the number of lactobacilli and bifidobacteria can negatively affect the formation of immune defense processes, predispose to the development of food allergies and inflammatory bowel diseases [11]. By controlling the functional state of the intestines of a child during the first months of life, it is possible to prevent the realization of a hereditary predisposition to allergies and the vaatopic phenotype [13].

The ability to protect a child predisposed to atopy from the influence of provoking factors can prevent its implementation into the phenotype of an allergic disease. Prevention of allergic diseases should be provided with a wide variety of measures and carried out at all stages of a person's life, starting from the prenatal period [17].

According to the World Health Organization (WHO), allergic diseases are currently one of the most significant problems, especially in pediatrics. In children, this pathology ranks second in prevalence. At the same time, there is an increase in the frequency of severe allergic reactions and an increasingly early onset of clinical manifestations. In this regard, timely and qualified diagnosis of allergic pathology in children is becoming the most urgent [17].

The growing prevalence of allergic diseases among children and adolescents remains one of the most important medical and social problems and causes a serious burden on the health budget of many countries of the world [14]. Allergic diseases negatively affect the physical and psychological state, social life, school performance and reduce the quality of life of both the patients themselves and their families [18].

However, despite the high incidence of morbidity, little attention is often paid to the problem of allergies: most pediatric patients either do not receive the necessary therapy at all, or are treated

from time to time, taking symptomatic drugs; in addition, self-medication is widespread.

In recent years, domestic and foreign researchers have published works on the clinical and immunological aspects of allergic diseases in children, the prevalence and intensity of these diseases in the child population, various methods of treatment, prevention of complications, treatment methods, prevention of complications and prevention of these pathologies.

The results obtained on the development of new methods for the diagnosis of allergic diseases in children, identified risk factors for the development of allergic diseases, including food allergies in children.

However, the research was mainly carried out in a hospital setting with an already developing pathological state, pre-pathological conditions were not taken into account, the relationship between risk factors for the development of allergies with living conditions and the ecological conditions of the region, there was practically no comprehensive, conceptual approach to the study of the formation and development of allergic diseases in rural areas. where there are no industrial enterprises and a sufficient allergenic background of the population, including children. In addition, there are few epidemiological, comprehensive studies to study the prevalence and intensity of detection of allergic diseases in rural areas of the republic. There are rare works on a comprehensive study of the prevalence of these diseases, clinical, immunobiological aspects of allergic diseases in children.

In this regard, research on a conceptual approach to a comprehensive clinical, immunological, medical and social study of allergic diseases among rural children, as well as the development of new criteria for early diagnosis, prognosis of their course and outcome is relevant and in demand.

Health disorders, in the medical sense, mean the onset, recurrence and progression of diseases. According to research by a number of authors, in the structure of the general morbidity in children aged 0 to 14 years, the 1st place is occupied by diseases of the respiratory system [50.1%], the 2nd - by diseases of the digestive system [6.5%], 3rd - diseases of the eye and its accessory apparatus (5%) [6, 7]. It should be emphasized that in the structure of the general morbidity of children, respiratory diseases consistently occupy the 1st place. According to L.I. Dziubich et al., Acute respiratory infections in children are the most common reason for parents to visit a children's clinic and account for about 90% of all childhood diseases [8].

The data from monitoring the physical development and health of children in preschool educational institutions indicate that only 17.3% of children can be considered absolutely healthy, 29.4% have functional abnormalities, 52.8% suffer from chronic diseases in the stage of compensation and 0.5% - chronic diseases in the stage of subcompensation [5].

In the structure of chronic diseases of older preschoolers, the first place belongs to diseases of the surgical sphere [3]. In second place are chronic diseases of the ENT organs [hypertrophy of the tonsils and adenoids of the II-III-degree, chronic rhinitis, sinusitis, otitis media]. In third place are diseases of the nervous system (episyndrome, hypertensive-hydrocephalic syndrome, the consequences of infantile cerebral palsy).

One of the most common pathologies among the child population is allergic diseases, which are a serious medical and social problem [4]. In terms of prevalence, severity of the disease, the level of socio-economic damage, the cost of treatment, allergic diseases were included in the first three groups of pathologies in the structure of human

diseases. According to modern scientific data, the prevalence of bronchial asthma, atopic dermatitis, and allergic rhinitis is noted in the structure of allergic diseases.

In a state of homeostasis, most hematopoietic progenitor cells are retained inside the bone marrow, only a small part of them continuously leaves its limits and enters the peripheral blood. Leukocytosis is a well-studied pathophysiological mechanism aimed at resisting infection, due to the gradual release of cellular elements from the bone marrow. Mobilization of CD34 + cells occurs when the balance between attachment and release of stem cells is imbalanced. If attachment is provided by the specific architectonics of the bone marrow, adhesion molecules and the production of chemokines that hold phenotypically immature cells in the bone marrow, then mobilization is the result of the action of “peripheral” chemokines (IL-8, etc.) and remodeling of the matrix and basement membrane by matrix enzymes (collagenase B or matrix metal proteinase-9 (MMP-9), etc.) [4]. The concept of circulating stem cells was first presented at the beginning of the 20th century by A.A. Maksimov, much later their presence in circulation was experimentally proved [15]. Data on the kinetics of CD34 + cells against the background of the inflammatory process are scarce and contradictory. For example, F. Mastrandrea et al. (2005) report that in patients with allergic inflammation, the content of circulating CD34 + cells is significantly higher than in patients with infectious fever and donors [16]. Other researchers have found that in patients with allergic diseases, both during remission and during an exacerbation, compared with healthy individuals comparable in gender, the level of CD34 lymphocytes in the peripheral blood does not change [1, 6, 8]. In patients with acute viral hepatitis B, C, B + C [5, 13], in children with septic complications of chemotherapy for

hemoblastosis [9] and acute intestinal infection [10], as well as against the background of systemic inflammation [16], a significant increase in phenotypically immature cells in circulation and a direct correlation with the level of C-reactive protein [11]. The lack of patient stratification by the severity of inflammation, the heterogeneity of the groups and the average level of evidence [there are no multicenter studies on this issue] do not allow judging the patterns of mobilization of phenotypically immature cells. Taking into account the universality of the mechanisms of inflammation that do not depend on the etiology of the pathological process, as well as relying on the results of our own research and the above data of other authors, we formulated a hypothesis that the severity of the inflammatory response correlates with the level of CD34 + cells in the bloodstream.

### Conclusion

The etiology of childhood health disorders is explained not only by genetic predisposition. The realization of hereditary burden is provoked by the constantly growing number of unfavorable environmental factors: increased environmental pollution by vehicles, poor-quality food, changes in immunity, under the influence of extensive vaccination programs, a decrease in the number of natural infections, etc.

### References:

1. Abdurakhmanov, Z. M., & Yemets, I. N. (2020). Long-term outcomes of aortic valve repair in children with congenital heart disease and their predictors. *Russian Journal of Cardiology*, 25(8), 131-138. doi:10.15829/1560-4071-2020-3971
2. Achilova, D. N., Amonov, R. A., Sharipova, L. K., Yomgurova, O. R., & Rustamov, B. B. (2021). *Clinical, Immunological and Medico-Social Aspects of Allergic Diseases*

- in Children. *Annals of the Romanian Society for Cell Biology*, 6736-6740.
3. Albitskii, V. Y. (2012). Social pediatrics: the history and development issues, the role of scientists of the scientific center of children's healthcare (Commencement speech of prof. V. Yu. Albitsky). *Rossiiskii Pediatricheskii Zhurnal*, 1, 4-10.
  4. Amrulloevich, G. S., & Norpulatovna, N. S. (2019). Maxillofacial anomalies in children with chronic tonsillitis and immunity factors, hypoxia and endogenous intoxication for the development and formation of pathology. *International Journal of Pharmaceutical Research*, 11(3), 1018-1026. doi:10.31838/ijpr/2019.11.03.103
  5. Erkara, I. P., Cingi, C., Ayranci, U., Gurbuz, K. M., Pehlivan, S., & Tokur, S. (2009). Skin prick test reactivity in allergic rhinitis patients to airborne pollens. *Environmental monitoring and assessment*, 151(1), 401-412.
  6. Imomjonovich, I. I., Fayzullayevich, S. S., & Erkinovich, N. J. -. (2021). Immunogenesis of kidney transplantation, maintenance of vital signs of transplanted kidney. *Annals of the Romanian Society for Cell Biology*, 25(3), 6794-6798. Retrieved from [www.scopus.com](http://www.scopus.com)
  7. Inoyatova, F. I., & Yusupalieva, G. A. (2016). Significance of complex echographic examinations in diagnosing chronic viral hepatitis in children. *Voprosy Detskoi Dietologii*, 14(1), 21-25. doi:10.20953/1727-5784-2016-1-21-25
  8. Gafurovich, V. U., & Kilichevna, K. M. (2020). The use of various techniques in the treatment of traumatic injuries of the oral mucosa in children. *European Journal of Molecular and Clinical Medicine*, 7(3), 3743-3748. Retrieved from [www.scopus.com](http://www.scopus.com)
  9. Khamidova, N. K., Mirzoeva, M. R., & Narzullaev, N. U. (2021). Clinical and immunological study of the effect of different types of therapy on the course of allergic rhinitis in children with hymenolepiasis. *Annals of the Romanian Society for Cell Biology*, 25(4), 1900-1908. Retrieved from [www.scopus.com](http://www.scopus.com)
  10. Kudratova, D. Sh, Ikhtiyarova, G.A., & Davlatov, S.S. (2021). Medical and social problems of the development of congenital malformations during a pandemic. *International Journal of Pharmaceutical Research*, 13(1), 756-760. doi: 10.31838/ijpr/2021.13.01.130
  11. Lauková, D. (2015). Medico-social aspects of patients with bronchial asthma. *Kontakt*, 17(2), e103-e115.
  12. Mariethoz, E., Lacroix, J. S., Moeschler, M. B., Hogendijk, S., Spertini, F., & Polla, B. S. (1999). Allergy and the Environment: A Meeting Report. *Reviews on environmental health*, 14(2), 63-78.
  13. Oripov, F., Blinova, S., Dekhkanov, T., & Davlatov, S. (2020). Development of immune structures of the leaning intestine of rabbits in early postnatal ontogenesis. *International Journal of Pharmaceutical Research*, 13(1), 299-301. doi:10.31838/ijpr/2021.13.01.042
  14. Teshayev, S. J., Khudoyberdiyev, D. K., & Davlatov, S. S. (2021). The impact of exogenous and endogenous factors on the stomach wall, macro-, microscopic anatomy of newborn white rats. *International Journal of Pharmaceutical Research*, 13(1), 679-682. doi:10.31838/ijpr/2021.13.01.101
  15. Vinogradov V.V. Stress and pathology. - Minsk: Belarusian. Science, 2007. - 351 p.
  16. Vishneva, E. A., Yuldashov, I. R., Nematova, K. G., & Yuldashov, S. I. (2020). Psychoemocial status in children with bronchial asthma. *Central Asian Journal of Pediatrics*, 2020(3), 106-116.