

Open Access Article

PECULIARITIES OF IMMUNE STATUS IN CHILDREN WITH CHRONIC BRONCHITIS

Nilufar Irgashevna Karimova

Republican Specialized Scientific and Practical Medical Center of Pediatrics, Ministry of Health of the Republic of Uzbekistan, Tashkent, Uzbekistan

E-mail: nilufar_karimova_00@mail.ru

Furkat Mukhitdinovich Shamsiev

Republican Specialized Scientific and Practical Medical Center of Pediatrics, Ministry of Health of the Republic of Uzbekistan, Tashkent, Uzbekistan

E-mail: furkat_shamsiev8388@mail.ru

Abstract

The aim of the study was to determine the state of the immune and cytokine status in chronic bronchitis in children. The results of the study showed that chronic bronchitis occurs in 10.4% of the total number of children with bronchopulmonary pathology and develops against the background of unfavorable perinatal factors, concomitant and past diseases affecting the development, course and outcome of the disease. The clinical picture and course of chronic bronchitis depend on the nature and degree of bronchial damage. The data obtained by us on the indicators of the immune status in different age groups in children with chronic bronchitis indicate suppression of indicators of cellular immunity and phagocytosis of neutrophils, activation of the humoral link of immunity, the most informative diagnostic markers of chronic diseases of the lower respiratory tract in children, manifested by overproduction of pro- and anti-inflammatory cytokines indicating chronic inflammation.

Keywords: chronic bronchitis, children, immune status, history, clinical manifestations

抽象的

该研究的目的是确定儿童慢性支气管炎的免疫状态和细胞因子状态。研究表明，慢性支气管炎发生在支气管肺病变儿童总数的10.4%，是在不利的围产期因素、伴随和既往疾病影响疾病发展、病程和结局的背景下发生的。慢性支气管炎的临床表现和病程取决于支气管损伤的性质和程度。我们获得的慢性支气管炎患儿不同年龄组免疫状态指标数据表明，细胞免疫指标和中性粒细胞吞噬功能受到抑制，免疫的体液环节被激活，是诊断慢性支气管炎最具信息量的标志物。儿童下呼吸道，表现为促炎和抗炎细胞因子的过度产生，表明慢性炎症。

关键词：慢性支气管炎，儿童，免疫状态，病史，临床表现

Introduction

Chronic lung diseases remain a serious public health problem in all countries of the world and

reach 10–40% among all pathologies of the bronchopulmonary system [1,2]. They lead to frequent, prolonged exacerbations, a decrease in

Received: April 12, 2021 / Revised: May 11, 2021 / Accepted: June 13, 2021 / Published: July 23, 2021

About the authors: Nilufar Irgashevna Karimova

Corresponding author- e-mail: nilufar_karimova_00@mail.ru

the quality of life of patients, disability at a young working age. The main place in the structure of chronic diseases of the lower respiratory tract belongs to chronic bronchitis [3,4,5]. Chronic bronchitis is a common disease, and the number of patients continues to increase, which, in all likelihood, is explained by the difficulties in identifying pathology due to the blurred clinical manifestations, the frivolous attitude of this contingent to the mild symptoms of the disease [6,7]. Immunological dysfunctions are an important factor contributing to the formation of chronic bronchitis in children. The failure of the immune system can be caused by genetic factors, immunodeficiency state, as well as unfavorable exogenous influence of the environment [8,9,10].

Purpose of the study. Determine the state of the immune and cytokine status in chronic bronchitis in children.

Materials And Methods

The study was carried out in 172 children with chronic bronchitis for the period 2020-2021, who were inpatient treatment at the pulmonology department. The comparison group consisted of 40 children with acute bronchitis. The control group consisted of 24 apparently healthy children of the same age. The diagnosis of chronic bronchitis was verified on the basis of complex clinical and instrumental studies: a carefully collected anamnesis, clinical, laboratory and instrumental (chest x-ray). The immunological study included the determination of CD3⁺, CD4⁺, CD8⁺, CD16⁺, CD20⁺ lymphocytes, the concentration of immunoglobulins of classes A, M, G, phagocytic activity of neutrophils, which was performed at the Institute of Human Immunology and Genomics.

Results

Analysis of anamnestic data in the examined children showed that the majority of children with chronic bronchitis were born from an unfavorable pregnancy with symptoms of toxicosis - 75.0% (129), 77.0% (132) of mothers of children with chronic bronchitis were diagnosed with anemia during pregnancy I and II degrees, which significantly ($P < 0.01$) differs from the indicators of the comparison group. Mothers of children with chronic bronchitis during this pregnancy suffered acute respiratory infections in 60.4% (104) cases, bronchitis 34.3% (59) had endocrine diseases 32.5% (56), which differs markedly from the indicators of the comparison group ($P < 0.01$). Most mothers had foci of chronic infection - diseases of the upper respiratory tract (rhinitis, tonsillitis) 69.7% (120), which differs markedly from the indicators of the comparison group ($P < 0.01$). It is known that the severity of chronic bronchitis in children is significantly influenced by an unfavorable premorbid background [11,12]. Analysis of the data showed that anemia of I-II degrees was determined in 94.7% (163) children, which differs from the indicators of the comparison group ($P < 0.001$); this, as is known, aggravates oxygen starvation in the child. Residual effects of rickets were detected in 54.0% (93), which complicates ventilation of the lungs.

The onset of the disease was acute in 29.0% (9) children with chronic bronchitis aged 3-6 years, in 31.2% (44) children aged 7-15 years, gradual in 71.0% (22) children in at the age of 3-6 years, in 68.8% (97) at the age of 7-15 years. The severity of the general condition was determined by the severity of clinical manifestations or complications. At the time of the present hospitalization, the general condition of the patients was severe in 28.5% (49) of children with chronic bronchitis, of moderate severity in

71.5% (123). Upon admission to the hospital, the main complaints of parents of children, patients were cough in 100.0% (172), shortness of breath - in 28.5% (49), fever - in 41.8% (72) children, decreased appetite - in 91.8% (158), lethargy - in 84.8% (146), pallor - in 92.4% (159). X-ray examination showed an increase and deformation of the bronchopulmonary pattern in 62.4% (88) children with chronic bronchitis, the process was more often localized in both lungs, in 167 (97.1%) patients there was an increase in the pulmonary pattern and in 85 (60.3%) deformation of the bronchopulmonary pattern was revealed in children. The carried out complex of immunological studies made it possible to assess the state of cellular and humoral immunity in chronic bronchitis. The results of the study of the immune status of the examined patients made it possible to establish a different severity of the immunodeficiency state (mainly due to the cellular link of immunity) with signs of tension in the humoral link of immunity and the multidirectional nature of immunological changes (Table 1). As can be seen from the

presented data, in patients with chronic bronchitis in the phase of exacerbation of the disease, the following deviations were revealed: a significant decrease in CD3⁺ lymphocytes, in relation to the comparison group (41.6±0.7% with 61.5±2.2% in children of the control group, P <0.001, in relation to the comparison group (50.2±2.5%, P <0.01). The indicators of children with chronic bronchitis exceeded those of children with acute bronchitis by 1.2 times. A significant decrease in CD3⁺ lymphocytes up to 36.8±0.6% at 61.5±2.2% in practically healthy children, P <0.001, in relation to the comparison group (50.2±2.5%; P <0.001). CD4⁺ lymphocytes belong to regulatory cells. Without them, transformation of CD20⁺ lymphocytes into plasma cells that form antibodies is impossible. A study of the number of CD4⁺ lymphocytes revealed their deep deficiency. The content of CD4⁺ lymphocytes was reduced to 28.9 281.0% in relation to the comparison group (P <0.01) in children with chronic bronchitis, which exceeded the acute bronchitis data by 1.3 times.

Table 1. Comparative analysis of indicators of cellular immunity in children with chronic bronchitis, (M ± m)

Indicators	Control group n = 24	Acute bronchitis n = 40 (I)	Chronic bronchitis n = 172 (II)	P
Leukocytes, abs.	6677,8±167,9	7048,5±125,9	6924,8±102,5	>0,05
Lymphocytes,%	33,5±0,5	35,2±0,6	38,2±0,5	<0,01
CD3 ⁺ - Lymphocytes,%	61,5±2,2	50,2±2,5	41,6±0,7	<0,01
CD4 ⁺ - Lymphocytes,%	39,1±2,1	37,9±1,3	28,9±1,0	<0,01
CD8 ⁺ - Lymphocytes,%	19,5±1,8	18,9±1,2	25,8±0,5	<0,001
(CD4 ⁺ /CD8 ⁺) immunoregulatory index	-2,0±0,2	2,0±0,2	1,1±0,06	<0,01
CD16 ⁺ - Lymphocytes,%	10,2±1,3	9,8±0,8	14,2±0,4	<0,01
Phagocytosis of neutrophils, %	58,5±2,3	52,1±1,1	39,8±1,2	<0,001

Note: P - reliability of differences between indicators of I and II groups of patients;

The number of CD8⁺ lymphocytes, which play an important role in antiviral immunity, in children with chronic bronchitis was increased by 1.3 times in comparison with the indicators of practically healthy children. There was a significant increase in CD8⁺ lymphocytes (25.8±0.5% at 19.5±1.8%; in children of the control group, P <0.001, in relation to the comparison group (18.9±1.2%; P <0.01). It was revealed a decrease in the immunoregulatory index in patients with chronic bronchitis by 1.8 times due to a more significant increase in the number of CD8⁺ cells compared to CD4⁺. There was a significant increase in natural killer cells - CD16⁺ lymphocytes in comparison with children

of the control group and amounted to 14.2 ± 0.4% (10.2 ± 1.3% in practically healthy children; P <0.001), in relation to the comparison group (P <0.05). There was a significant decrease in neutrophil phagocytosis to 39.8±1.2% in relation to the comparison group (52.1±1.1%, P <0.01). The change in neutrophil phagocytosis was more pronounced in the group of children with chronic bronchitis compared with acute bronchitis (P <0.001). of these children, there was also a tendency to an increase in CD20⁺ lymphocytes (32.4±0.7% at 16.4±0.5% in children of the control group, P <0.001, in relation to the comparison group (P <0.01) (table 2).

Table 2. Comparative analysis of indicators of humoral immunity in the examined children, (M ± m)

Indicators	Control group (n = 24)	Acute bronchitis n = 40 (I)	Chronical bronchitis n = 172 (II)	P
CD20 ⁺ -Lymphocytes,%	16,4±0,5	18,1±0,8	32,4±0,7	<0,001
IgG, mg/%	938,3±17,6	908,7±18,4	1360,5±36,5	<0,001
IgA, mg /%	107,9±3,6	104,6±2,8	149,7±5,8	<0,001
IgM, mg /%	90,7±2,8	102,5±3,4	126,8±2,5	<0,01

Note: P - reliability of differences between indicators of I and II groups of patients;

The activation of the humoral link of immunity was evidenced by the increased concentration of immunoglobulin IgM in patients; their increase was 1.4 times, on average, up to 126.8 ± 2.5 mg /%, in relation to the comparison group by 1.2 times (102.5±3, 4mg /% P <0.01). The IgG content in the group of patients with chronic bronchitis showed an increase in this indicator by 1.5 times compared with the data of practically healthy children and averaged 1360.5 ± 36.5 mg

/% (P <0.001), in relation to the comparison group it was increased 1.5 times (908.7 ± 18.4 mg /%, P <0.001). In the examined children with chronic bronchitis, an increase in the IgA level was also noted by 1.4 times compared with the indicators of practically healthy children and on average was 149.7 ± 5.8 mg /% (P <0.001), it was increased in relation to the comparison group 1.4 times (104.6±2.8 mg /%, P <0.001).

Conclusion

Thus, summarizing the results of the study, we can conclude that chronic bronchitis occurs in 10.4% of the total number of children with bronchopulmonary pathology and develops against the background of unfavorable perinatal factors, concomitant and transferred diseases affecting the development, course and outcome of the disease. The clinical picture and course of chronic bronchitis depend on the nature and degree of bronchial damage. The data obtained by us on the indicators of the immune status in different age groups in children with chronic bronchitis indicate suppression of indicators of cellular immunity and phagocytosis of neutrophils, activation of the humoral link of immunity, the most informative diagnostic markers of chronic diseases of the lower respiratory tract in children, manifested by overproduction of pro- and anti-inflammatory cytokines indicating chronic inflammation.

References

1. Samsygina G.A. Chronic bronchitis in children: modern vision of the problem // *Pediatrics*. -2016. -№3.-C.55-59.
2. Korolev A.V., Blokhin B.M. Respiratory diseases in children - M., ID "Medpraktika-M", 2017.- 616s.
3. Baranov A.A., Khaitov R.M. Allergology and Immunology. Clinical guidelines for pediatricians. Publishing house: Union of Pediatricians of Russia. -2018.-492s.
4. Chuchalin A.G., Avdeev S.N., Aisanov Z.R., Belevsky A.S., Leshchenko I.V., Meshcheryakova N.N., Ovcharenko S.I., Shmelev E.I. Federal clinical guidelines for the diagnosis and treatment of chronic obstructive pulmonary disease // *Pulmonology*. - 2020. -№3. -S.15-36.
5. Kim D.H, Yoo I.Y. Development of a questionnaire to measure resilience in children with chronic diseases // *J Korean Acad Nurs*. 2019 Apr; 40 (2): 236-46.
6. Chuchalin A.G., Khaltaev N., Antonov N.S., Galkin D.V., Manakov L.G., Antonini P., Murphy M., Solodovnikov A.G., Bousquet J., Pereira M.H., Demko I.V. Chronic respiratory diseases and risk factors in 12 regions of the Russian Federation. *Int J Chron Obstruct Pulmon Dis* 2014; 9: 963-974.
7. The chronic bronchitic phenotype of COPD: an analysis of the COPD Gene Study [Text] / V. Kim, M. K. Han, G. B. Vance [et al.] // *Chest*. - 2017. - Vol. 140, No. 3. - P. 626-633.
8. Sereda E.V., Selimzyanova L.R., Kustova O. V., Lukina O. F., Lokhmatov M.M. Chronic bronchitis in congenital and hereditary respiratory diseases in children. Modern diagnostic technologies and therapeutic tactics // *Russian Pediatric Journal*. - 2018.-№4.-C.36-40.
9. Shamsiev A.M., Yusupov Sh. A., Yuldashev B.A., Mukhamadieva L.A. The state of the immune status in children with chronic bronchitis // *Pediatric Bulletin of the Southern Urals*. - 2017.- No. 1.-P.84-89.
10. Barnes P.J. Immunology of asthma and chronic obstructive pulmonary disease // *Nat. Immunology. Rev*. - 2018. - No. 8. - P. 183-192.
11. Barnes P.J., Shapiro S.D., Pauwels R.A. Chronic obstructive pulmonary disease: molecular and cellular mechanisms // *Eur. Respir. J*. - 2018. - Vol. 22. - P. 672–688.
12. Oppenheim J., Feldman M. Cytokine reference. -London: Academic Press, 2000.-P. 356-412;